



Contents lists available at SciVerse ScienceDirect

# International Journal for Parasitology

journal homepage: [www.elsevier.com/locate/ijpara](http://www.elsevier.com/locate/ijpara)



## How effective is preening against mobile ectoparasites? An experimental test with pigeons and hippoboscid flies

Jessica L. Waite\*, Autumn R. Henry, Dale H. Clayton

Department of Biology, University of Utah, Salt Lake City, UT, USA

### ARTICLE INFO

**Article history:**  
Received 29 December 2011  
Received in revised form 7 March 2012  
Accepted 8 March 2012  
Available online xxxxx

**Keywords:**  
Grooming  
Behaviour  
Defence  
*Columba livia*  
*Pseudolynchia canariensis*  
Vector

### ABSTRACT

Birds combat ectoparasites with many defences but the first line of defence is grooming behaviour, which includes preening with the bill and scratching with the feet. Preening has been shown to be very effective against ectoparasites. However, most tests have been with feather lice, which are relatively slow moving. Less is known about the effectiveness of preening as a defence against more mobile and evasive ectoparasites such as hippoboscid flies. Hippoboscids, which feed on blood, have direct effects on the host such as anaemia, as well as indirect effects as vectors of pathogens. Hence, effective defence against hippoboscid flies is important. We used captive Rock Pigeons (*Columba livia*) to test whether preening behaviour helps to control pigeon flies (*Pseudolynchia canariensis*). We found that pigeons responded to fly infestation by preening twice as much as pigeons without flies. Preening birds killed twice as many flies over the course of our week-long experiment as birds with impaired preening; however, preening did not kill all of the flies. We also tested the role of the bill overhang, which is critical for effective preening against feather lice, by experimentally removing the overhang and re-measuring the effectiveness of preening against flies. Birds without overhangs were as effective at controlling flies as were birds with overhangs. Overall, we found that preening is effective against mobile hippoboscid flies, yet it does not eliminate them. We discuss the potential impact of preening on the transmission dynamics of blood parasites vectored by hippoboscid flies.

© 2012 Published by Elsevier Ltd. on behalf of Australian Society for Parasitology Inc.

### 1. Introduction

Birds are infested with a variety of ectoparasites including lice, mites, ticks, fleas and flies, all of which have the capacity to decrease host fitness (Atkinson et al., 2008; Møller et al., 2009). Birds combat ectoparasites with defences ranging from anti-parasite behaviour (Hart, 1992, 1997) to immune defences (Wikel, 1996; Owen et al., 2010). Grooming behaviour, which includes preening with the bill and scratching with the feet, is the first line of defence against ectoparasites (Clayton et al., 2010). Preening is an energetically expensive activity (Goldstein, 1988; Croll and McLaren, 1993); furthermore, the time and energy devoted to preening detracts from other behaviours such as feeding and vigilance (Redpath, 1988). Therefore, in order to be effective against ectoparasites while limiting its energetic cost, preening should be an inducible defence (Tollrian and Harvell, 1999). The importance of preening is illustrated by recent work demonstrating that features of bill morphology, such as the upper mandibular overhang, appear to have evolved specifically to enhance the effectiveness of preening for parasite control (Clayton and Walther, 2001; Clayton et al., 2005).

Nearly all of the work on the effectiveness of preening has been done with feather lice (Phthiraptera: Ischnocera), which are slow moving and therefore relatively easy targets for preening birds (Marshall, 1981; Atkinson et al., 2008). The effectiveness of preening for controlling more mobile ectoparasites such as fleas and hippoboscid flies has not, to our knowledge, been tested. Preening may also play a role in shaping vector ecology and the evolution of pathogens transmitted by ectoparasites.

The goal of our study was to test the effectiveness of preening against hippoboscid flies, which are mobile parasites of birds and mammals. Avian hippoboscid flies are dorso-ventrally flattened and very agile at slipping between the feathers. As described by Rothschild and Clay (1952): “They have... an extremely efficient method of moving among feathers – darting and scuttling about at a remarkable speed – and are extremely difficult to catch on a living bird.” Hippoboscids may also be capable of avoiding preening by using “refugia” such as the vent region of the bird or behind the bases of the legs (Waite, personal observation).

Hippoboscid flies are a diverse group of parasites. More than 200 species are recognised, 75% of which parasitise birds belonging to 18 orders; the rest parasitise mammals (Lloyd, 2002; Lehane, 2005). Most species of bird flies are winged and capable of flight between individual hosts (Harbison et al., 2009; Harbison and Clayton, 2011). They spend most of their time on the body of the

\* Corresponding author. Tel.: +1 801 585 9742; fax: +1 801 581 4668.  
E-mail address: [jessi.waite@gmail.com](mailto:jessi.waite@gmail.com) (J.L. Waite).

bird, where they feed on blood several times a day (Coatney, 1931). Hippoboscid feeding can cause anaemia (Jones, 1985), emaciation (Lloyd, 2002) and slow nestling development (Bishopp, 1929). Parents of hippoboscid-infested nestlings have lower reproductive success (Bize et al., 2004). Hippoboscid flies also transmit blood parasites that can have negative effects on birds, including malaria (Sol et al., 2003), trypanosomes (Baker, 1967) and possibly viruses such as West Nile (Farajollahi et al., 2005). In short, hippoboscids pose both direct and indirect threats to the health and fitness of their hosts.

To test the effectiveness of preening against hippoboscid flies, we used wild caught Rock Pigeons (*Columba livia*) that we experimentally infested with the pigeon fly *Pseudolynchia canariensis* (Diptera: Hippoboscidae). We conducted two separate experiments. The first experiment addressed two questions: (i) do Rock Pigeons infested with flies increase the amount of time they spend preening and (ii) is preening effective in killing flies? The second experiment addressed a third question: is the bill overhang important in the effectiveness of preening for fly control?

## 2. Materials and methods

### 2.1. Experiment 1: preening and flies

Twenty-four Rock Pigeons were caught using walk-in traps in Salt Lake City, Utah, USA. The birds were transported to the University of Utah animal facility, where they were individually housed in wire mesh cages (30 × 30 × 56 cm) suspended over newspaper-lined trays. Each cage/tray was completely enclosed within a fly-proof net, which prevented flies from moving between birds in different cages. Birds were given ad libitum food, water and grit and kept in a 12-h light/dark cycle. They were maintained in captivity for at least 6 months at low humidity prior to the experiment, which killed feather lice and their eggs that were present on the birds when they were captured (Harbison et al., 2008). Any flies present on pigeons when they were captured would have died during the 6 month period because the life span of pigeon flies is only 2–3 months (Fahmy et al., 1977). Since pigeons trapped in Salt Lake City do not usually have other ectoparasites, the birds were ectoparasite-free at the start of our experiment. Prior to the start of the experiment, birds were carefully examined to confirm that they did not, in fact, have any ectoparasites.

We blocked the 24 birds using two factors: (i) location trapped and (ii) time in captivity; we then randomly assigned birds to one of three treatments, with eight birds per treatment. All birds were sexed and weighed. Birds in the first two treatments were then infested with 20 flies each (10 male flies, 10 female flies), which is the maximum number recorded from wild pigeons (mean = 5.07 flies; Stekhoven et al., 1954). Flies used to infest birds were cultured from wild caught stock on pigeons kept for this purpose in another room. The third group of eight birds was not infested with flies.

Flies were removed from culture birds using CO<sub>2</sub> (Moyer et al., 2002). They were sexed under a microscope at 25× before putting them on experimental birds. Half of the birds (Chosen at random) in each of the two fly-infested treatments had plastic attachments fitted to their bill to impair their ability to preen. The attachments are small C-shaped pieces of plastic that, when fitted in the nares of a pigeon, create a 1.0–3.0 mm gap between the mandibles. This gap prevents the full occlusion of the bill needed for effective preening (Clayton et al., 2005). The attachments are harmless; they do not impair feeding or alter the amount of time that birds attempt to preen (Clayton and Tompkins, 1995; Koop et al., 2011).

To address our first question whether pigeons preen more when they are infested with flies, we compared the behaviour of birds with normal (unimpaired) preening with and without flies. Preen-

ing behaviour was quantified using instantaneous scan sampling between 13:00 and 16:00 h (Altmann, 1974). Preening was defined as touching the plumage with the bill (Clayton and Cotgreave, 1994). Birds were observed at 6 s intervals (Clayton, 1990) for 30 observations per bird per day, for 5 days following infestation. We calculated the proportion of time that birds spent preening.

To address our second question whether preening is effective in killing flies, we compared the number of flies killed by birds with impaired preening with flies killed by birds with normal preening. The experiment lasted 1 week, after which one of the authors (Waite) removed dead flies from the bottom of each cage; food and water dishes were also checked for dead flies. Another author (Henry) re-examined all cages to ensure that nothing was overlooked. Damage to flies was observed and recorded under a microscope at 25×. Flies were scored as preening-damaged if the head, thorax, abdomen or at least one wing was crushed or missing, or if at least three legs were missing. We calculated the proportion of flies with preening-damage out of the total number of dead flies recovered for each host after 1 week.

### 2.2. Experiment 2: bill overhang

Another 12 wild-caught (individually caged) pigeons were used for this experiment. Birds were again blocked by location trapped and time in captivity. Half of the birds, chosen at random, had their bill overhang trimmed away with a dremel tool. The other half was sham trimmed, i.e. they were handled but no part of the bill was removed (Fig. 1). The trimming method, which is fully described in Clayton et al. (2005), does not harm the birds in any way. One week after trimming (or sham trimming) all birds were sexed and weighed, and then each bird was infested with 20 flies (10 males, 10 females). Preening behaviour and fly mortality were quantified as in Experiment 1.

### 2.3. Statistical analysis

Statistical analyses were performed in Prism® v. 5.0b (GraphPad Software, Inc.). Data were analysed using Mann-Whitney U Tests




Fig. 1. Rock Pigeon bill showing upper mandibular overhang before (A) and after (B) removal of the overhang. The overhang grows back after several weeks. Figure reproduced from Clayton et al. (2005).

184 for comparisons between two groups. ANOVAs were used for compar-  
 185 comparisons among three groups. The sex ratio of pigeon hosts in each  
 186 experiment was compared using a Chi-square or Fisher's Exact test,  
 187 as appropriate. Values are presented as mean ± S.E. Results were  
 188 considered significant at  $P \leq 0.05$ .

189 **3. Results**

190 Sex and body mass of hosts did not differ significantly by treat-  
 191 ment in either experiment (Experiment 1: sex, Chi-square test,  
 192  $P = 0.77$ ; mass, ANOVA,  $F_{2,21} = 1.47$ ,  $P = 0.25$ ; Experiment 2: sex,  
 193 Fisher's Exact test,  $P = 1.00$ ; mass, **Mann-Whitney**  $U = 12.5$ ,  
 194  $P = 0.42$ ).

195 **3.1. Experiment 1: preening and flies**

196 Birds infested with flies preened more than twice as much as  
 197 birds without flies; birds with flies preened  $23.49 \pm 3.96\%$  of the  
 198 time observed, whereas birds without flies preened  $11.21 \pm 2.11\%$   
 199 of the time observed; (Fig. 2). The difference in preening rates be-  
 200 tween the two groups was statistically significant (**Mann-Whitney**  
 201  $U = 10.5$ ,  $P = 0.03$ ).

202 Birds with normal preening killed twice as many flies as birds  
 203 with impaired preening; birds with normal preening killed  
 204  $43.75 \pm 5.41\%$  of flies, compared with  $21.88 \pm 5.74\%$  of flies killed  
 205 by birds with impaired preening (Fig. 3A). The difference in the  
 206 number of flies killed was statistically significant ( $U = 11.0$ ,  
 207  $P = 0.03$ ).

208 Birds with normal preening also damaged a significantly greater  
 209 proportion of dead flies than did birds with impaired preening  
 210 (Fig. 3B; **Mann-Whitney**  $U = 7.0$ ,  $P = 0.01$ ). Of the dead flies recov-  
 211 ered from normally preening birds,  $44.6 \pm 0.06\%$  were damaged,  
 212 while only  $16.6 \pm 0.13\%$  of flies recovered from birds with impaired  
 213 preening were damaged.

214 **3.2. Experiment 2: bill overhang**

215 Removal of the bill overhang had no significant effect on preening-  
 216 time; birds without overhangs preened  $12.96 \pm 1.08\%$  of the  
 217 time observed, while birds with overhangs preened  $16.81 \pm 3.90\%$   
 218 of the time observed (**Mann-Whitney**  $U = 13.0$ ,  $P = 0.47$ ). Birds with  
 219 overhangs did not kill significantly more flies than birds with no  
 220 overhang; birds with overhangs killed  $50.83 \pm 11.93\%$  of flies, com-  
 221 pared with  $45.00 \pm 11.76\%$  of flies killed by birds with no overhang  
 222 (Fig. 4; **Mann-Whitney**  $U = 15.0$ ,  $P = 0.69$ ). Thus, the bill overhang  
 223 was not a factor in the efficiency with which preening killed flies.




Fig. 2. Proportion of time that birds with and without flies spent preening.




Fig. 3. Effect of preening and an example of preening damage. (A) Proportion of flies killed by birds with normal versus impaired preening. (B) Example of intact versus preening-damaged flies.




Fig. 4. Proportion of flies that were dead in cages of birds with and without bill overhangs.

4. Discussion

We examined the effectiveness of preening against mobile ectoparasitic flies. Pigeons experimentally infested with flies preened twice as much as pigeons without flies (Fig. 2). Preening also

proved to be effective against flies (Fig. 3A); we recovered twice as many dead flies from the cages of birds that could preen, compared with those that could not preen. Pigeons were able to catch and crush flies (Fig. 3B), even though the flies are extremely adept at moving quickly and evasively through the feathers (Rothschild and Clay, 1952).

Removal of the bill overhang did not decrease the efficiency of preening significantly (Fig. 4). Clayton et al. (2005) showed that lice are crushed when birds preen by the mortar-and-pestle action of the tip of the lower mandible moving against the upper mandibular overhang. Although the overhang is essential for controlling feather lice, our results show that it is not needed when preening flies, presumably because the flies are much larger and softer-bodied than lice. Although preening proved to be an effective defence against flies, it did not eliminate all of them over the course of our week-long experiment. Only one of 40 birds in the two experiments cleared itself completely of flies.

Preening may have the added benefit of helping to protect birds from pathogens for which the flies are vectors. In principle, preening can prevent transmission of pathogens if it kills infected vectors before they have an opportunity to bite the host. The fly *P. canariensis* is a known vector of the blood parasites *Haemoproteus columbae* and *Trypanosoma hanna* (Fahmy et al., 1977; Mandal, 1991). Waite (unpublished data) recently showed that pigeons exposed to just five flies for 3 days can become infected with *H. columbae*. In our study, only an average of 50% of flies placed on pigeons were killed during the week-long experiment (Fig. 3A). Thus, even birds with relatively efficient preening may remain at risk of acquiring blood parasites. If preening irritates flies, encouraging them to move between hosts, then preening might even have the effect of increasing pathogen transmission (Hodgson et al., 2001). It would be very interesting to measure the impact of preening on pathogen transmission by hippoboscids among birds in a population.

We found that pigeons infested with flies doubled the amount of time that they spent preening compared with controls (without flies) and compared with the typical rates of preening for other pigeons and doves (Clayton, 1990; Koop et al., 2011). One might predict that experimental birds would spend even more time preening, given that they did not completely remove their infestations in most cases. However, research on the cost of preening shows that it is energetically expensive. When birds preen, their metabolic rates increase by as much as 200% (Wooley, 1978; Croll and McLaren, 1993). The energetic cost of preening might explain why preening is an inducible defence against hippoboscid flies. Additional indirect costs of preening include the time taken away from courtship behaviour, foraging and predator surveillance (Redpath, 1988). Thus, in addition to the direct impact of hippoboscid flies on host fitness, flies may have indirect effects mediated by the energetic and time related costs of preening. Indeed, there may well be a trade-off between the indirect cost of preening and the more direct costs of fly infestation.

## Acknowledgements

All work was performed with the University of Utah IACUC, USA, approval (Protocol #08-08004). We thank Sung Ki Hong for assistance with data collection and animal care, and Kari Smith for assistance in maintaining the fly culture. We thank Jennifer Koop for help with behavioural data collection methods and Sarah Bush for discussion and help with graphics. We are grateful to Franz Goller, Jael Malenke and Lesley Chesson for comments on the manuscript. We thank three anonymous reviewers, whose comments improved the manuscript. We thank the Royal Society, UK for permission to reproduce Fig. 1. Funding was provided by Sigma Xi, USA and the American Ornithologists Union, USA to

J.L.W., the University of Utah Undergraduate Bioscience Research Program to A.R.H. and the National Science Foundation, USA DEB-0816877 to D.H.C.

## References

- Altmann, J., 1974. Observational study of behavior: sampling methods. *Behaviour* 49, 227–267.
- Atkinson, C.T., Thomas, N.J., Hunter, D.B., 2008. *Parasitic Diseases of Wild Birds*. Wiley-Blackwell, Iowa.
- Baker, J.R., 1967. A review of the role played by the Hippoboscidae (Diptera) as vectors of endoparasites. *J. Parasitol.* 53, 412–418.
- Bishopp, F.C., 1929. The pigeon fly – an important pest of pigeons in the United States. *J. Econ. Entomol.* 22, 947–987.
- Bize, P., Roulin, A., Tella, J.L., Bersier, L.-F., Richner, H., 2004. Additive effects of ectoparasites over reproductive attempts in the long-lived alpine swift. *J. Anim. Ecol.* 73, 1080–1088.
- Clayton, D.H., 1990. Mate choice in experimentally parasitized rock doves: lousy males lose. *Integr. Comp. Biol.* 30, 251–262.
- Clayton, D.H., Cotgreave, P., 1994. Relationship of bill morphology to grooming behaviour in birds. *Anim. Behav.* 47, 195–201.
- Clayton, D.H., Tompkins, D.M., 1995. Comparative effects of mites and lice on the reproductive success of rock doves (*Columba livia*). *Parasitology* 110, 195–206.
- Clayton, D.H., Walther, B., 2001. Influence of host ecology and morphology on the diversity of Neotropical bird lice. *Oikos* 94, 455–467.
- Clayton, D.H., Moyer, B.R., Bush, S.E., Jones, T.G., Gardiner, D.W., Rhodes, B.B., Goller, F., 2005. Adaptive significance of avian beak morphology for ectoparasite control. *Proc. Biol. Sci.* 272, 811–817.
- Clayton, D.H., Koop, J., Harbison, C., Moyer, B., Bush, S., 2010. How birds combat ectoparasites. *Open Ornithol. J.* 3, 41–71.
- Coatney, G., 1931. On the biology of the pigeon fly, *Pseudolynchia maura* Bigot (Diptera, Hippoboscidae). *Parasitology* 23, 525–532.
- Croll, D.A., McLaren, E., 1993. Diving metabolism and thermoregulation in common and thick-billed murrelets. *J. Comp. Physiol. B* 163, 160–166.
- Fahmy, M., Mandour, A., Arafat, M., Makhloof, L., 1977. Bionomics and natural infection of *Pseudolynchia canariensis* with *Haemoproteus* and *Trypanosoma* in Assuit Area, Upper Egypt. *J. Egypt. Soc. Parasitol.* 7, 19–24.
- Farajollahi, A., Crans, W.J., Nickerson, D., Bryant, P., Wolf, B., Glaser, A., Andreadis, T.G., 2005. Detection of West Nile virus RNA from the louse fly *Icosta americana* (Diptera: Hippoboscidae). *J. Am. Mosq. Control Assoc.* 21, 474–476.
- Goldstein, D., 1988. Estimates of daily energy expenditure in birds: the time-energy budget as an integrator of laboratory and field studies. *Am. Zool.* 28, 829–844.
- Harbison, C.W., Bush, S., Malenke, J., 2008. Comparative transmission dynamics of competing parasite species. *Ecology* 89, 3186–3194.
- Harbison, C.W., Jacobsen, M.V., Clayton, D.H., 2009. A hitchhiker's guide to parasite transmission: the phoretic behaviour of feather lice. *Int. J. Parasitol.* 39, 569–575.
- Harbison, C.W., Clayton, D.H., 2011. Community interactions govern host-switching with implications for host–parasite coevolutionary history. *Proc. Natl. Acad. Sci. USA* 108, 9525–9529.
- Hart, B.L., 1992. Behavioral adaptations to parasites: an ethological approach. *J. Parasitol.* 78, 256–265.
- Hart, B.L., 1997. Behavioural defence. In: Clayton, D.H., Moore, J. (Eds.), *Host-Parasite Evolution: General Principles and Avian Models*. Oxford University Press, Oxford, pp. 59–77.
- Hodgson, J., Spielman, A., Komar, N., Krahnforst, C., Wallace, G., Pollack, R., 2001. Interrupted blood-feeding by *Culiseta melanura* (Diptera: Culicidae) on European starlings. *J. Med. Entomol.* 38, 59–66.
- Jones, C., 1985. Heavy hippoboscid infestations on buzzards. *Br. Birds* 78, 592.
- Koop, J.A.H., Huber, S.K., Clayton, D.H., 2011. Does sunlight enhance the effectiveness of avian preening for ectoparasite control? *J. Parasitol.* 98, 46–48.
- Lehane, M.J., 2005. *The Biology of Blood-sucking in Insects*, 2nd ed. Cambridge University Press, New York.
- Lloyd, J., 2002. Louse flies, keds, and related flies (*Hippoboscoidea*). In: Mullen, G., Durden, L. (Eds.), *Medical and Veterinary Entomology*. Academic Press, Boston, pp. 349–362.
- Mandal, F.B., 1991. A preliminary report on the incidence of blood parasites in pigeon *Columba livia* and pigeon fly, *Pseudolynchia canariensis* (Macquart). *Indian J. Anim. Health.* 30, 29–32.
- Marshall, A.G., 1981. *The Ecology of Ectoparasitic Insects*. Academic Press Inc., New York.
- Moyer, B., Gardiner, D., Clayton, D.H., 2002. Impact of feather molt on ectoparasites: looks can be deceiving. *Oecologia* 131, 203–210.
- Møller, A., Arriero, E., Lobato, E., Merino, S., 2009. A meta-analysis of parasite virulence in nestling birds. *Biol. Rev. Camb. Philos. Soc.* 84, 567–588.
- Owen, J.P., Nelson, A.C., Clayton, D.H., 2010. Ecological immunology of bird-ectoparasite systems. *Trends Parasitol.* 26, 530–539.
- Redpath, S., 1988. Vigilance levels in preening dunlin *Calidris alpina*. *Ibis* 130, 555–557.
- Rothschild, M., Clay, T., 1952. *Fleas, flukes and cuckoos*. Collins, London.
- Sol, D., Jovani, R., Torres, J., 2003. Parasite mediated mortality and host immune response explain age-related differences in blood parasitism in birds. *Oecologia* 135, 542–547.

373 Stekhoven Jr., J., Silva, I., San Roman, P., 1954. Zur biologie der Taubenausfliege  
374 (Diptera, Pupipara). Z. Parasitenkunde 16, 388–406.  
375 Tollrian, R., Harvell, C. (Eds.), 1999. The Ecology and Evolution of Inducible Defenses.  
376 Princeton University Press, Princeton, NJ.

Wikel, S.K. (Ed.), 1996. The immunology of host–ectoparasitic arthropod  
relationships. CAB International, Guildford.  
Wooley Jr., J., 1978. Energy costs of activity and daily energy expenditure in the  
black duck. J. Wildl. Manage. 42, 739–745.

377  
378  
379  
380  
381

UNCORRECTED PROOF